Review

Innate Immunity and Cell Death in **Alzheimer's Disease**

SangJoon Lee¹, Hyun-Jeong Cho², and Jin-Hyeob Ryu^{3,4}

Abstract



American Society or Neurochemistry

ASN Neuro Volume 13: 1-16 © The Author(s) 2021 Article reuse guidelines: sagepub.com/iournals-permissions DOI: 10.1177/17590914211051908 journals.sagepub.com/home/asn

(S)SAGE

The innate immune system plays key roles in controlling Alzheimer's disease (AD), while secreting cytokines to eliminate pathogens and regulating brain homeostasis. Recent research in the field of AD has shown that the innate immune-sensing ability of pattern recognition receptors on brain-resident macrophages, known as microglia, initiates neuroinflammation, AB accumulation, neuronal loss, and memory decline in patients with AD. Advancements in understanding the role of innate immunity in AD have laid a strong foundation to elucidate AD pathology and devise therapeutic strategies for AD in the future. In this review, we highlight the present understanding of innate immune responses, inflammasome activation, inflammatory cell death pathways, and cytokine secretion in AD. We also discuss how the AD pathology influences these biological processes.

Keywords

alzheimer's disease, amyloid- β , tau, innate immunity, inflammasome, NLRP3, MxA, ASC speck, IL-1 β , IL-18, caspase-1, pyroptosis, apoptosis, necroptosis, neuroinflammation

Received July 14, 2021; Revised September 13, 2021; Accepted for publication September 21, 2021

Introduction

Alzheimer's disease (AD), the most common dementing illness, is characterized by progressive memory decline and cognitive dysfunction. AD affects more than 44 million people worldwide, with a new patient diagnosed every 4 s. The number of patients with dementia, including AD, is expected to double every 20 years and reach 115 million by 2050, placing huge financial burdens on individuals and society.

The three pathological hallmarks of AD are deposition of extracellular β -amyloid (A β), intraneuronal aggregation of neurofibrillary tangles composed of the microtubule-associated protein tau, and neuronal loss called neurodegeneration (Nelson et al., 2012) (Figure 1). In mammalian hosts, $A\beta$ and tau deposits as danger-associated molecular patterns (DAMPs) are recognized and cleared by multiple patternrecognition receptors (PRRs) including Toll-like receptors (TLRs), retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs), and nucleotide-binding oligomerization domain-like receptor family proteins (NLRs) (de Rivero Vaccari et al., 2014; Heneka et al., 2013; Ising et al., 2019; Tahara et al., 2006; Venegas et al., 2017).

In response to DAMPs, some PRRs can assemble large multiprotein complexes known as inflammasomes (Place &

Kanneganti, 2018). Upon assembly, the inflammasomes induce membrane pore formation and proinflammatory cytokine processing, leading to a form of inflammatory cell death known as pyroptosis (He et al., 2015; Man et al., 2017; Shi et al., 2015). Innate immune signaling and inflammasome activation are key protective responses against AD (Heneka et al., 2013, 2014; Venegas et al., 2017). However, their activation must be tightly regulated, as excessive activation can lead to neuroinflammation and brain damage. Balancing the

³BIORCHESTRA Co. Ltd., 17, Techno 4-ro, Yuseong-gu, Daejeon, Republic of Korea

⁴BIORCHESTRA Co. Ltd., 245 Main St, Cambridge, MA 02142, USA

Corresponding Authors:

Hyun-Jeong Cho, PhD, Department of Biomedical Laboratory Science, College of Medical Science, Konyang University, Daejeon, South Korea. Tel: +82-42-600-8433 Email: hjcho@konyang.ac.kr

Jin-Hyeob Ryu, PhD, B615~619, 17, Techno 4-ro, Yuseong-gu, Daejeon, Korea.

Email: branden.ryu@biorchestra.com

¹Department of Infection Biology, Faculty of Medicine, University of Tsukuba, Tsukuba, lapan

²Department of Biomedical Laboratory Science, College of Medical Science, Konyang University, Daejeon, South Korea

Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution- \odot \odot NonCommercial 4.0 License (https://creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access page (https://us.sagepub.com/enus/nam/open-access-at-sage).



Figure 1. Aggregation of A β and tau. The A β precursor APP undergoes processing by proteases α -, β -, and γ -secretase. A β monomers can aggregate into A β oligomers, protofibrils, fibrils, and ultimately plaques, which are among the hallmarks of AD pathology. Loss of microtubule binding leads to elevated levels of cytosolic tau, thereby increasing the potential for tau-tau interactions. Aggregation of hyperphosphorylated tau protein leads to the assembly of neurofibrillary tangles followed by progressive cytoskeletal changes, disruption of axonal transport, and AD pathology.

host innate immune response has been considered in potential therapeutic approaches for AD, but this balance must be finely tuned to reduce excessive neuroinflammation, while clearing A β and tau deposits.

In this review, we discuss innate immune recognition, inflammasome activation, programmed cell death, and



Tau spreading in neuron

Figure 2. A β and tau pathology in AD. Extracellular A β and tau contribute to enhanced bioactivity of microglia, which induces the secretion of proinflammatory cytokines including IL-1 β and IL-18. Activated microglia further take up $A\beta$ and tau and ultimately induce impaired immune responses in microglia and sterile inflammation. Several forms of tau contribute to neuron-to-neuron spreading, uptake, and aggregation of tau, thereby leading to tau-induced toxicity.

cytokine release that occur in response to AD, as well as how innate immune responses regulate the disease to resist development of AD pathology.

Pathogenesis of Alzheimer's Disease

AD is the most common type of dementia and neurodegenerative disease involving neuroinflammation, neuronal loss, and memory decline (Heneka et al., 2015; Nussbaum & Ellis, 2003). A β deposition can begin in the brain decades before the onset of clinical symptoms and stimulate tau-mediated AD pathogenesis such as tau hyperphosphorylation, tau aggregation, and triggering neuron-to-neuron spread (Figure 2).

As AB accumulation can induce the formation of neurofibrillary tangles that propagate to produce mild cognitive impairment and dementia, Aß production has been extensively characterized (Jack et al., 2010). Aß is a cleavage product of amyloid precursor protein (APP). APP expressed in neurons and glial cells has several physiological roles in mediating cell-to-cell adhesion for neuronal signaling and neurotransmitter release. APP cleavage by proteolytic secretases can generate amyloid peptides, including a peptide of 37–49 amino acids, A β , and released into the extracellular space in the brain (Chen et al., 2017). APP processing can follow the following two pathways: amyloidogenic and nonamyloidogenic. The amyloidogenic pathway involves a cleavage event induced by β -secretase such as BACE1, which releases soluble APP_β. The APP C-terminal fragment can then be cleaved by γ -secretases such as presentiin (PS) 1 and 2 at one of the several sites varying from 40 to 44

amino acid residues to generate neurotoxic A β peptides. Among Aßs of various lengths, soluble $A\beta_{1-42}$ is a major component of plaque formation and initiates Aß aggregation due to its hydrophobic C-terminus, which reduces the flexibility of transformation from α -helical to β -sheet structure. The non-amyloidogenic pathway is mediated by α -secretases such as ADAM 9, 10, and 17, and γ -secretase catalyzes further cleavages to produce non-toxic and possibly beneficial soluble APPa fragments with important neuroprotective functions (Chasseigneaux & Allinquant, 2012; Chen et al., 2017). The A β_{1-42} peptide can induce the production of higher-order assemblies of A β , wherein the monomers of A β peptide accumulate by misfolding into propagating conformations to freely diffuse through the brain (Chen et al., 2017). Aβ oligomers act as seeds to form insoluble A β aggregates know as plaques (Katzmarski et al., 2020). Aß plaques disrupt cell-to-cell communication, lead to neuronal apoptosis, and activate immune cells such as microglia, thereby triggering inflammation and brain tissue damage (Heneka et al., 2015). Activated microglia further take up A β and tau and ultimately induce impaired immune responses in microglia and sterile inflammation (Busche & Hyman, 2020) (Figure 2).

Tau is a type II microtubule-associated protein that is highly expressed in neurons and moderately expressed in oligodendrocytes and astrocytes in the mammalian brain. Under normal condition, tau mainly modulates the stability of microtubules in axons by direct interaction with tubulin. Tau also has synaptic functions regulated by post-translational modifications (PTMs), including phosphorylation and proteolytic cleavage (Wesseling et al., 2020). However, abnormal PTMs induce tau hyperphosphorylation, which causes it to dissociate from microtubules and to enhance tau aggregation (self-assembly) (Alonso et al., 2018). Therefore, hyperphosphorylation of tau substantially impairs the stability of microtubules in nerve cells, which is involved in the pathogenesis of AD. The hyperphosphorylated tau is released by exosomal processes, which can be detected in the cerebrospinal fluid and blood of patients with AD (Fiandaca et al., 2015; Jia et al., 2019), with the amounts detected correlating with cognitive impairment. Although exosomal tau is a potential biomarker for AD, it is necessary to consider how to achieve high sensitivity with diagnostic tools early in the onset of AD. Extracellular misfolded tau released from pathological cells via exosomes can enter naive cells such as neurons and microglia, leading to tau-induced toxicity and convert monomeric physiological tau via endocytosis (Hardy & Selkoe, 2002; He et al., 2018; LaFerla et al., 1995; Selkoe & Hardy, 2016) (Figure 2).

Innate Immune Recognition in Alzheimer's Disease

As resident immune effector cells of the central nervous system, microglia play a crucial role in regulating brain homeostasis and mediating innate immune responses in AD (Clayton et al., 2017; Hanisch & Kettenmann, 2007). Microglia sense a variety of microbial molecules known as pathogen-associated molecular patterns (PAMPs) as well as host-derived DAMPs via PRRs, including TLRs, RLRs, and NLRs (Kigerl et al., 2014) (Table 1). Microglia associated with AD are commonly in an activated state with upregulated expression of PRRs. PRRs drive signal transduction pathways that induce an inflammatory response and secretion of pro-inflammatory cytokines, including type I interferons (IFNs), thereby leading to the microglia-associated clearance of debris via phagocytosis in AD (Heneka et al., 2014; McDonough et al., 2017). Although microglia are integral for the phagocytosis and degradation of AB, their chronic activation can provoke impaired neuroinflammatory responses, with the potential to induce $A\beta$ production and neuronal distress (Hansen et al., 2018; Hemonnot et al., 2019; Sarlus & Heneka, 2017).

TLR- and RLR-Mediated Alzheimer's Disease Recognition

TLR- and RLR-mediated signaling leads to the secretion of type I IFNs and proinflammatory cytokines. TLRs are membranebound PRRs expressed in the microglia of mice, rats, and humans, and their activation can be beneficial or detrimental to the host (Bsibsi et al., 2002; Olson & Miller, 2004; Zhang et al., 2013). In microglia, the activation of TLR2 together with CD14, a coreceptor of TLR4, induces an immune response associated with fibrillar A β phagocytosis (Reed-Geaghan et al., 2009). Moreover, TLR2 and TLR4 activation in microglia enhances the production of A β peptides with 42 amino acids $(A\beta_{1-42})$, which are key pathogenic species in AD (Chen et al., 2006). Some studies have shown that the inhibition of TLR2 activation attenuates glial cell reactivity, leading to reduced AB deposits and improved cognitive function in APP/ PS1 double transgenic mice (McDonald et al., 2016). Furthermore, Tlr2/4-deficient mice are protected from neurocognitive and behavior impairment after immunization with A β_{1-42} peptide (Vollmar et al., 2010). Tlr2^{-/-} microglia induce proinflammatory cytokines, including tumor necrosis factor (TNF)- α , interleukin (IL)-1 β , and IL-6 and enhance A β clearance (Jana et al., 2008; Liu et al., 2012).

APP/PS1 transgenic mice of an AD model deficient in TLR4 exhibit reduced microglial activation, reduced cognitive function, and increased A β deposition (Song et al., 2011). In a transgenic mouse model carrying a mutation in tau (P301S), mild and chronic stimulation of TLR4 with lipopolysaccharide reduced the level of cerebral phosphorylated tau proteins and improved memory (Qin et al., 2016). In contrast, the activation of TLR4 reduced the clearance of microglial A β_{1-42} peptides by modulating the activity of the scavenger receptor CD36 (Li et al., 2015). Pronounced TLR4 expression in APP transgenic mice and increased TLR4 expression in the brains of patients with AD are

Sensor	Extracellular Aβ deposits	Intracellular neurofibrillary tangles (pTau)	Inflammatory cytokines	Cognition function	Reference
TLR2	Increase	N/A	Increase (TNF-α, IL-1β, IL-8)	Impair	(Liu et al., 2012)
	Increase	N/A	N/A	Impair	(McDonald et al., 2016)
TLR4	Decrease	N/A	Increase (IL-Iβ)	Improve	(Song et al., 2011)
	N/A	Decrease	Increase (TNF-α, IL-1β, IL-8)	Improve	(Qin et al., 2016)
	Increase	N/A	Increase (IL-6, TNF-α)	Impair (AD patient)	(Walter et al., 2007)
	Increase	N/A	Increase (TNF-α, IL-1β, IL-6)	Impair	(Balducci et al., 2017)
RIG-I	Increase	N/A	N/A	Impair	(de Rivero Vaccari et al., 2014)
NLRP3	Increase	N/A	Increase (IL-Iβ)	Impair	(Heneka et al., 2013)
	N/A	Increase	Increase (IL-I β)	Impair	(Ising et al., 2019)
	Increase	Increase	Increase (IL-I β)	N/A	(Stancu et al., 2019)
	Increase	N/A	N/A	N/A	(Yin et al., 2018)
	Increase	N/A	Increase (TNF-α, IL-1β, IL-6)	Impair	(Lonnemann et al., 2020)
AIM2	Increase	N/A	No significant (IL-Iβ)	No significant	(Wu et al., 2017)
Extracellular ASC	Increase	N/A	Increase (IL-I β)	N/A	(Friker et al., 2020)
speck	Increase	N/A	N/A	Impair	(Venegas et al., 2017)

Table 1. Effects and Functions of Innate Immune Sensor in AD.

associated with A β plaques (Walter et al., 2007). The activation of TLR4 is necessary for glial cell activation, which resulted in memory impairment in a mouse model of AD (Balducci et al., 2017). A recent study suggested that alterations in microglial TLR4 signaling contribute to AD pathogenesis in APP/PS1 double transgenic mice (Go et al., 2016). Collectively, the activation of TLR2 and TLR4 in the initial stages of AD has beneficial effects in A β clearance, whereas chronic activation contributes to A β aggregation.

RIG-I is a cytosolic PRR that canonically senses 5'-triphosphate double-stranded RNA. Although one study revealed that RIG-I was elevated in the temporal cortex and plasma of patients with mild cognitive impairment (de Rivero Vaccari et al., 2014), the detailed molecular mechanism of how RIG-I is involved in AD pathology remains unclear. Thus, further studies are required to identify the *in vivo* dominant innate sensor in AD.

A β complexed with nucleic acids triggers TLR- and RLR-derived type I IFN responses in microglia, resulting in complement-mediated synapse destruction in AD (Roy et al., 2020). Therefore, IFNs constitute a pivotal element within the neuroinflammatory network of AD and critically contribute to neuropathogenic processes.

NLRP3 Inflammasomes in AD

Innate immune cells in the brain play major roles in cytokine production and inflammatory signaling. Among the

proinflammatory cytokines, IL-1ß and IL-18 are the key mediators of the inflammatory response; increased levels of these proteins correlate with the severity of AD in patients (Forlenza et al., 2009; King et al., 2018; Ng et al., 2018; Ojala et al., 2009). The release of IL-1β and IL-18 requires proteolytic maturation of pro-IL-1ß and pro-IL-18. This process is mediated by inflammasome formation and protease caspase-1 activation (Man et al., 2017). Inflammasome sensors can recognize PAMPs and/or DAMPs produced during pathogen infection and cellular instability, respectively (Kanneganti, 2010). Among the sensors, the most wellcharacterized sensor is the NLR family pyrin domaincontaining 3 (NLRP3), which has been implicated in several diseases such as autoinflammatory diseases, obesity, colitis, and pathogen-mediated diseases including influenza A viruses and corona viruses (Gurung & Kanneganti, 2016; Kanneganti et al., 2006a, 2006b; Karki et al., 2015; Lee et al., 2020; Lee & Ryu, 2021; Stienstra et al., 2011; Thomas et al., 2009; Zaki et al., 2010).

The NLRP3 inflammasome has emerged as a trigger of AD pathogenesis. The mRNA and protein expressions of NLRP3 are upregulated in the monocytes of patients with AD (Saresella et al., 2016). Loss of the NLRP3 inflammasome (NLRP3/caspase-1) in APP/PS1 mice protects against long-term potentiation deficits and attenuates spatial memory deficits and the A β burden (Heneka et al., 2013).

Furthermore, exogenous-aggregated tau activates the NLRP3 inflammasome in the microglia (Stancu et al.,

2019), and fibrillar A β -containing brain homogenates induce tau pathology in an NLRP3-dependent manner (Ising et al., 2019). Consistently, NLRP3 inflammasome inhibitor (MCC950, blocks NLRP3-ASC-induced IL-1 β secretion and ASC oligomerization) reduces phosphorylation of tau and accumulation of A β in the hippocampus induced by exogenous tau seeding in TauP301S transgenic mice (Stancu et al., 2019). Other NLRP3 inflammasome inhibitors, such as the drug JC-124 and fenamate non-steroidal anti-inflammatory drugs, exert beneficial effects in mouse models of AD (Daniels et al., 2016; Yin et al., 2018). Collectively, these findings suggest that the NLRP3 inflammasome contributes to AD pathology.

MxA Inflammasomes in AD

Besides NLRP3, MxA also acts as an inflammasome sensor in respiratory epithelial cells (Lee et al., 2018, 2019). The Mx genes, which are present in almost all vertebrates, are interferon-regulated genes whose expression is upregulated following IFN production, which leads to an antiviral response, particularly against RNA viruses (Haller et al., 2015). Following influenza A virus infection, MxA recognizes the influenza virus A nucleoprotein, which then interacts with the inflammasome adapter protein and apoptosis-associated speck-like protein containing a CARD (ASC), and leads to caspase 1 activation and IL-1 β and IL-18 production (Lee et al., 2018, 2019).

It has been reported that MxA polymorphisms are associated with risk and age-at-onset in AD and accelerate cognitive decline in patients with AD (Ma et al., 2012). Another study showed that MxA is found in senile plaques of the AD brain



Figure 3. Extracellular ASC specks in AD. Activation of microglia by a TLR agonist leads to transcriptional induction of genes encoding components of the NLRP3 inflammasome and pro-IL-1 β . Upon A β and tau uptake, lysosomal damage leads to the assembly of the NLRP3 inflammasome, which eventually induces the release of IL-1 β and oligomerization of ASC to form ASC specks. Extracellular ASC specks interact with A β aggregates in the extracellular space to promote their deposition as plaques. Extracellular ASC specks can be engulfed by neighboring microglia and contribute to the exacerbation of neuroinflammation via further production of IL-1 β .

and the presence of MxA in reactive microglia could contribute to AD pathology (Yamada et al., 1994). Given that IL-1 β and IL-18 are associated with the severity of AD in patients (Forlenza et al., 2009; King et al., 2018; Ng et al., 2018; Ojala et al., 2009), it will be interesting to reveal whether MxA inflammasome is required for AD exacerbation.

Extracellular ASC Specks in AD

Inflammasome sensors, such as NLRP3, are intracellular proteins expressed in macrophages and are activated by a wide variety of DAMPs, such as alum, silica, uric acid, and amyloid- β , following their engulfment by macrophages. Activated NLRP3 facilitates ASC oligomerization to form large, intracellular macromolecular aggregates known as ASC specks. It has been reported that ASC specks are released into the extracellular space, where ASC propagates inflammatory responses via prion-like transmission mediated by engulfment in neighboring macrophages (Baroja-Mazo et al., 2014; Franklin et al., 2014). In AD pathology, in the extracellular space, ASC specks bind to $A\beta_{1-42}$ to accelerate its oligomerization, indicating the cross-seeding activity of ASCs in A β aggregation, which boosts its toxicity in microglia (Friker et al., 2020; Venegas et al., 2017) (Figure 3). Recently, it has been reported that the extracellular ASC specks are engulfed by Arf6-deficient macrophages, and IL-1 β production is reduced in Arf6^{-/-} macrophages compared with that in wild-type macrophages (Lee et al., 2021a). Although detailed molecular mechanisms of how intracellular ASC specks are released into the extracellular space and the role of these molecules in neuroinflammation remain largely unknown, the extracellular ASC specks may induce Arf6 dependent-secondary inflammasomes in neighboring microglia and AD pathology.

Programmed Cell Death and Proinflammatory Cytokines in Alzheimer's Disease

Cell death plays an important role in limiting disease progression. However, inflammatory cell death also results in the release of proinflammatory cytokines and cellular contents, including PAMPs and DAMPs, which can induce severe inflammation (Bergsbaken et al., 2009; Pasparakis & Vandenabeele, 2015) (Figure 4). Therefore, cell death is considered a double-edged sword during disease progression.

Pyroptosis in Alzheimer's Disease

Pyroptosis is a form of inflammatory cell death mediated by inflammasomes and gasdermin (Shi et al., 2015). Inflammasome assembly leads to the activation of inflammatory caspases [caspase-1 (human and mouse), -4 (human), -5 (human), or -11 (mouse)], which proteolytically cleave and



Figure 4. Programmed cell death pathways in AD pathology. In AD, $A\beta$ and tau deposition act as danger-associated molecular patterns (DAMPs) and stimulate inflammasome assembly, resulting in the activation of caspase-1. Activated caspase-1 cleaves pro–IL-1 β , pro–IL-1 β , and gasdermin D (GSDMD). The N-terminal fragment of GSDMD may then oligomerize within membranes to form membrane pores and execute pyroptosis. A β and tau deposition initiates a signaling cascade mediated by caspase-8 and –9 activation. Caspase-8 and –9 induce the activation of caspase-3 to drive apoptosis. AD exacerbation by A β and tau deposition initiates necroptosis, a receptor-interacting serine/ threonine-protein kinase 1 (RIPK1) and RIPK3 complex-dependent form of inflammatory cell death that depends on the activation of the protein mixed lineage kinase domain-like pseudokinase (MLKL) to form channels in the membrane.

release the N-terminal fragment of gasdermin D (GSDMD) to form pores in the plasma membrane (He et al., 2015; Man et al., 2017; Shi et al., 2015). Caspase-1-dependent GSDMD cleavage leads to the release of IL-1 β and IL-18 (Man et al., 2017).

Recombinant $A\beta_{1-42}$ can induce caspase-1 activation, GSDMD cleavage, and pyroptosis in mouse cortical neurons (Han et al., 2020; Tan et al., 2014). NLRP3 inflammasome and caspase-1 activation, along with IL-1 β release, have also been observed when fibrillar A β is phagocytosed by microglia (Halle et al., 2008). Furthermore, recombinant tau protein can activate the NLRP3 inflammasome and induce IL-1 β release from microglia (Stancu et al., 2019). A recent study revealed that microglia isolated from Tau22 mice, which express human tau mutations involved in frontotemporal dementia, stimulate pyroptosis, as evidenced by NLRP3 inflammasome and caspase-1 activation and IL-1 β release (Ising et al., 2019).

Necroptosis in Alzheimer's Disease

Necroptosis is a form of inflammatory cell death mediated by mixed-lineage kinase domain-like pseudokinase (MLKL). MLKL is oligomerized by the phosphorylation of receptorinteracting serine/threonine-protein kinase 1 (RIPK1) and RIPK3; the oligomerized MLKL is translocated to the plasma membrane and forms channels (Dondelinger et al., 2014).

In the temporal gyrus of human patients with AD, RIPK1 and MLKL show robust expression, and necroptosis is associated with reduced brain weight of patients with AD (Caccamo et al., 2017). Furthermore, necroptosis exacerbates cognitive deficits in APP/PS1 transgenic mice (Caccamo et al., 2017). Another study revealed that APP/PS1 mice treated with pharmacological inhibitors of RIPK1 or *Ripk1^{D138N/D138N}* mice exhibit reductions in A β burden, inflammatory cytokine levels, and memory deficits (Ofengeim et al., 2017).

Apoptosis in Alzheimer's Disease

Apoptosis is executed by caspase-3 and -7 following the activation of upstream initiator caspases including caspase-8/10 or -9. Pyroptosis and necroptosis are inflammatory and lytic cell death processes in which cytokines are released to induce inflammation and alert immune cells, whereas apoptosis has been historically considered to be immunologically silent. However, recent studies have suggested that apoptosis is not always immunologically silent, as crosstalk occurs between apoptotic proteins and the molecules executing lytic cell death (Lee et al., 2020; Place et al., 2021). The apoptotic protein caspase-3 has been reported to activate gasdermin E to induce lytic cell death (Rogers et al., 2019). Caspase-8 activates GSDMD under specific conditions (Lee et al., 2020; Place et al., 2021).

In the context of AD, APP is a substrate for caspase-3mediated cleavage, which contributes to A β plaque formation, synaptic loss, and behavioral changes associated with AD (Gervais et al., 1999). Other studies have reported that the activation of caspase-3, -8, and -9 further induces A β plaque formation and AD progression (Rohn et al., 2001, 2002; Stadelmann et al., 1999; Su et al., 2001). Collectively, these studies suggest that the apoptotic pathways may also contribute to AD progression.

Recent studies have found extensive crosstalk between programmed cell death pathways, establishing the concept of PANoptosis (pyroptosis, apoptosis, necroptosis) (Gurung et al., 2014; Kuriakose et al., 2016; Lamkanfi et al., 2008; Lee et al., 2020, 2021b; Malireddi et al., 2010, 2021; Place et al., 2021). Although inflammasome activation is primarily associated with gasdermin D (GSDMD)-mediated pyroptosis, there is emerging evidence of the contribution of inflammasome components in driving PANoptosis (Christgen et al., 2020; Gurung et al., 2014; Kesavardhana et al., 2020; Kuriakose et al., 2016; Lee et al., 2020, 2021b; Malireddi et al., 2020, 2021; Place et al., 2021; Zheng et al., 2020).

The AIM2 innate immune sensor, which is activated by double-stranded DNA, has been implicated in AD, that is, *Aim2* deficiency reduces A β deposition and microglial activation (Wu et al., 2017). Recently, it has been reported that AIM2 regulates the innate immune sensors pyrin and ZBP1 to drive inflammatory signaling and PANoptosis (Lee et al., 2021b). It will be interesting to reveal whether AIM2 PANoptosis exerts beneficial effects in AD.

Inflammatory Cytokines

Microglia-associated PRRs drive signal transduction pathways that induce inflammatory cell death pathways and secretion of pro-inflammatory cytokines, including type I IFNs, thereby leading to the microglia-associated clearance of debris via phagocytosis in AD (Heneka et al., 2014; McDonough et al., 2017).

However, over activation of inflammatory cell death pathways can lead to critical brain damage and severe neuroinflammation (Bergsbaken et al., 2009; Pasparakis & Vandenabeele, 2015). Inflammatory cell death leads to the release of proinflammatory cytokines and chemokines, which may contribute to AD pathology.

Indeed, A β -stimulated human monocytes release chemokines such as IL-8, monocyte chemoattractant protein-1 (MCP-1), macrophage inflammatory protein 1 α (MIP-1 α), and MIP-1 β *in vitro*. Microglia cultured from rapid autopsies of patients with AD and nondemented patients revealed increased expression of IL-8, MCP-1, and MIP-1 α after experimental exposure to A β (Xia & Hyman, 1999).

Many proinflammatory cytokines have been shown to be produced by neurons or microglia and the levels of IL-1 α , IL-1β, IL-6, TNF-α, granulocyte-macrophage colony-stimulating factor, and IFN- α are increased in AD brain tissue (Friker et al., 2020; Ising et al., 2019; Rubio-Perez & Morillas-Ruiz, 2012; Stancu et al., 2019; Venegas et al., 2017). Proinflammatory cytokines such as IL-18, IL-1 β , IL-6, and TNF- α are released by astrocytes and microglia in response to A β plaques or following tau hyperphosphorylation (Forloni et al., 1997; Wang et al., 2015). Recent studies revealed that the combination of TNF- α and IFN- γ induces inflammatory cell death, tissue damage, and cytokine shock syndromes (Karki & Kanneganti, 2021; Karki et al., 2021). Furthermore, treating with neutralizing antibodies against TNF- α and IFN- γ protected mice during SARS-CoV-2 infection, sepsis, hemophagocytic lymphohistiocytosis, and cytokine shock (Karki & Kanneganti, 2021; Karki et al., 2021). Therefore, it is important to identify the innate immune processes contributing to cytokine release and potential associations between cytokine signaling and inflammatory cell death in AD pathology.

Innate Immune Signals Regulate Adaptive Immunity

The innate immune and adaptive immune systems interact with each other. Although innate immune responses control the progression of AD, adaptive immune responses also contribute to restraining AD pathogenesis. B and T lymphocytes are the main cellular components of adaptive immunity that are responsible for antigen-specific antibody secretion and cell-mediated immunity, respectively. The adaptive immune response plays a key role in the development of adequate control against toxic molecules, such as misfolded tau and A β (Anderson et al., 2014; Cantrell, 2015). The triple transgenic (3xTg-AD) mouse model shows tau and A β accumulation in the brain increasing with age and changes in their immune system. A recent study has shown that CD4⁺/CD8⁺ lymphocyte ratio in the blood of 3xTg-AD mice increased compared with that of WT mice (St-Amour et al., 2014), suggesting a universal deficit in the adaptive immune response; the finding is consistent with abnormal lymphocyte populations in patients with AD (Larbi et al., 2009; Pellicanò et al., 2012; Pirttilä et al., 1992; Saresella et al., 2011; Schindowski et al., 2006; Speciale et al., 2007).

Innate immune sensing by PRRs instructs adaptive immunity (Palm & Medzhitov, 2009). Therefore, innate and adaptive immune systems function cooperatively, and crosstalk between peripheral and central immunity such as cytokine and chemokine signaling likely plays an important albeit understudied role in AD. Recent studies have shown that peripheral neutrophils and T-regulatory cells profoundly affect AD pathogenesis (Baruch et al., 2015; Zenaro et al.,

Agent	Target/ Mechanism ^a	Mechanism of action	Clinical trial phaseª	Reference
Atuzaginstat (COR388)	Inflammation/ Infection	Small molecule; bacterial protease inhibitor targets gingipain produced by <i>Porphyromonas gingivalis</i> ; reduces neuroinflammation	III	(Haditsch et al., 2020)
Azeliragon (TTP488)	Amyloid/ inflammation	Small molecule inhibitor; RAGE antagonist; reduces Aβ transport into the brain; mitigates toxic effects of oligomers and reduces inflammation	111	(Lue et al., 2001)
NE3107 (HE30286)	Inflammation	MAPK-1/3 inhibitor; reduces proinflammatory NFκB activation	III	(Ahlem et al., 2009)
AL002 (anti-TREM2)	Inflammation	Immunotherapy; monoclonal antibody; targets microglial TREM2 receptors; promotes microglial clearance of Aβ and reduces neurotoxicity	II	(Wang et al., 2020)
ALZT-OPI (cromolyn + ibuprofen)	Inflammation	Small molecule and combination therapy (cromolyn; approved anti-asthma drug, ibuprofen; approved anti-inflammatory drug); reduces aggregation of Aβ and induces neuroprotective microglial activation	II	(Brazier et al., 2017; Zhang et al., 2018)
Daratumumab (anti-CD38)	Inflammation/ Immunity	Immunotherapy (FDA-approved for the treatment of multiple myeloma); monoclonal antibody; targets CD38 on glia cells; regulates microglial activity	II	(Blacher et al., 2015; Guerreiro et al., 2020)
Dasatinib + quercetin	Inflammation/ Immunity	Senolytic therapy; tyrosine kinase inhibitor (dasatinib) and flavonoid (quercetin, nutritional supplement); reduces senescent cells and tau aggregation	II	(Zhang et al., 2019)
GB301	Inflammation/ Immunity	Cell therapy drug; autologous regulatory T cells; reduces neuroinflammation	II	(Dansokho et al., 2016; Plascencia-Villa and Perry 2020)

Table 2.	Target and Immunit	y Mechanism of	Therapeutic Agents	in Clinical	Trials for	Alzheimer's Disease
----------	--------------------	----------------	--------------------	-------------	------------	---------------------

Dasatinib + quercetin	Inflammation/ Immunity	CD38 on glia cells; regulates microglial activity Senolytic therapy; tyrosine kinase inhibitor (dasatinib) and flavonoid (quercetin, nutritional supplement); reduces senescent cells and tau aggregation	II	(Zhang et al., 2019)
GB301	Inflammation/ Immunity	Cell therapy drug; autologous regulatory T cells; reduces neuroinflammation	II	(Dansokho et al., 2016; Plascencia-Villa and Perry, 2020)
Lenalidomide	Inflammation/ Immunity	FDA-approved cancer drug; reduces inflammatory cytokines; regulates innate and adaptive immune responses	II	(Decourt et al., 2020)
Montelukast (MK0476)	Inflammation	Small molecule; cysteinyl leukotriene type I (cysLT-1) receptor antagonist; affects inflammatory processes, neuronal injury, blood-brain-barrier integrity, and Aβ protein accumulation	II	(Lai et al., 2014; Morin et al., 2014)
Pepinemab (VX15)	Inflammation	Immunotherapy; monoclonal antibody inhibitor of semaphoring 4D (SEMA4D); reduces inflammatory cytokine release	II	(LaGanke et al., 2017)
AL003	Inflammation	Immunotherapy; monoclonal antibody targeting SIGLEC-3 (CD33); reactivates microglia and immune cells in the brain; improves microglial clearance of toxic proteins	I	(Estus et al., 2019)
Edicotinib	Inflammation	Small molecule; CSF-1R antagonist; blocks microglial proliferation and production of cytokines (IL1 β and TNF α)	I	(Mancuso et al., 2019)
XPro I 595	Inflammation	Protein biologic; soluble TNF α inhibitor; reduces neuroinflammation	Ι	(MacPherson et al., 2017)

^aTarget/mechanism and clinical trial phases are based on ClinicalTrials.gov (2021).

2015). A recent study revealed A β plaque formation, neuroinflammation, and microglial activation in 5XFAD mice (Marsh et al., 2016). Additionally, the loss of IgG-producing B cells impairs microglial phagocytosis, thereby inducing A β deposition and worsening AD pathology (Marsh et al., 2016).

Adaptive immunity in AD pathology has not been studied using genetic deletion of key PRRs. The use of knockout mice lacking key innate sensors in specific immune cell subsets will reveal the cell type-specific requirements for these sensors in instructing various aspects of adaptive immunity in AD pathology.

Therapies for Alzheimer Disease

Currently, there is no effective therapy for AD except aducanumab; a new therapy for early AD was approved by the FDA in 2021. Aducanumab is a human monoclonal antibody that selectively targets aggregated A β (Sevigny et al., 2016). Given the increasing incidence of AD globally, effective treatment strategies for AD are urgently needed. Over the last two decades, several drugs that decrease A β production or increase A β clearance in the brain have been identified. Although A β plaques are the neuropathological hallmark of AD, they are poorly correlated with AD severity and cognitive dysfunction (Arriagada et al., 1992; Giannakopoulos et al., 2003). Patients with AD in whom brain A β plaques were cleared by anti-A β immunotherapy showed no cognitive benefit (Holmes et al., 2008).

One possible target for intervention is neuroinflammation. Indeed, neuroinflammatory biomarkers shed light on therapeutic target candidates such as soluble TREM2, RAGE, CD38, CD33, and TNF- α (Table 2). Several studies have consistently identified a link between the inflammasome and AD, and microglia play a central role in linking inflammation with neurodegeneration (Friker et al., 2020; Heneka et al., 2013; Salter & Stevens, 2017; Venegas et al., 2017). In contrast, an efficient adaptive immune system may prevent AD pathogenesis by modulating microglial function (Marsh et al., 2016). Further studies are needed to understand the role of the innate immune system and microglia to explore whether immune-based therapies can be designed for AD.

Tau-targeted therapies are another active area of investigation. As tau pathology correlates better with cognitive impairments than A β lesions, targeting tau is expected to be more effective than A β clearance once clinical symptoms are evident (Congdon & Sigurdsson, 2018). Unfortunately, most initial anti-tau therapies based on the inhibition of kinases or tau aggregation or on stabilization of microtubules have been discontinued due to their lack of efficacy and toxicity (Congdon & Sigurdsson, 2018). Currently, most tautargeted therapies in clinical trials are immunotherapies, which have shown promise in numerous pre-clinical studies (Congdon & Sigurdsson, 2018).

Conclusions

Research in the past two decades have substantially expanded our understanding of innate immune responses in AD pathology. Emerging evidence suggests that aberrant microglial function in innate immune responses and cell death are inextricably linked to AD pathology. Therefore, innate immune responses and cell death must be finely controlled to reduce excessive neuroinflammation during the clearance of deposited A β and tau. Nevertheless, the following questions remain unanswered. What is the dominant innate immune sensor-derived signaling pathway that controls neuroinflammation and clearance of $A\beta$ and tau deposition? What types of cell death are involved in neuroinflammation in AD? How can the innate immune response be balanced to reduce excessive neuroinflammation while retaining its support for the adaptive immune response? This association among inflammasome, inflammatory cell death, and neuroinflammation in AD is being actively pursued by numerous studies, and exploring the roles of these factors will help identify new drug targets for combating this devastating neurodegenerative disease.

Acknowledgments

We apologize to our colleagues in the field whose work could not be cited because of space limitations.

Author Contributions

S.L., H.C., and J.R. outlined the manuscript. S.L., H.C., and J.R. wrote the manuscript. S.L., H.C., and J.R. critically revised and approved the final version of the manuscript.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Ministry of Education, Culture, Sports, Science, and Technology of Japan (grant number 19K16665).

ORCID iD

SangJoon Lee D https://orcid.org/0000-0001-7500-782X

References

- Ahlem, C., Auci, D., Mangano, K., Reading, C., Frincke, J., Stickney, D., & Nicoletti, F. (2009). HE3286: A novel synthetic steroid as an oral treatment for autoimmune disease. *Annals of the New York Academy of Sciences*, 1173, 781–790. https://doi.org/ 10.1111/j.1749-6632.2009.04798.x
- Alonso, A. D., Cohen, L. S., Corbo, C., Morozova, V., Elidrissi, A., Phillips, G., & Kleiman, F. E. (2018). Hyperphosphorylation of tau associates with changes in its function beyond microtubule

stability. *Frontiers in Cellular Neuroscience*, *12*, 338. https://doi.org/10.3389/fncel.2018.00338

- Anderson, K. M., Olson, K. E., Estes, K. A., Flanagan, K., Gendelman, H. E., & Mosley, R. L. (2014). Dual destructive and protective roles of adaptive immunity in neurodegenerative disorders. *Translational Neurodegeneration*, 3(1), 25. https:// doi.org/10.1186/2047-9158-3-25
- Arriagada, P. V., Growdon, J. H., Hedley-Whyte, E. T., & Hyman, B. T. (1992). Neurofibrillary tangles but not senile plaques parallel duration and severity of Alzheimer's disease. *Neurology*, 42(3 Pt 1), 631–639. https://doi.org/10.1212/WNL. 42.3.631
- Balducci, C., Frasca, A., Zotti, M., La Vitola, P., Mhillaj, E., Grigoli, E., Iacobellis, M., Grandi, F., Messa, M., Colombo, L., Molteni, M., Trabace, L., Rossetti, C., Salmona, M., & Forloni, G. (2017). Toll-like receptor 4-dependent glial cell activation mediates the impairment in memory establishment induced by β-amyloid oligomers in an acute mouse model of Alzheimer's disease. *Brain Behavior and Immunity*, *60*, 188–197. https://doi.org/10.1016/j. bbi.2016.10.012
- Baroja-Mazo, A., Martín-Sánchez, F., Gomez, A. I., Martínez, C. M., Amores-Iniesta, J., Compan, V., Barberà-Cremades, M., Yagüe, J., Ruiz-Ortiz, E., Antón, J., Buján, S., Couillin, I., Brough, D., Arostegui, J. I., & Pelegrín, P. (2014). The NLRP3 inflammasome is released as a particulate danger signal that amplifies the inflammatory response. *Nature Immunology*, 15(8), 738–748. https://doi.org/10.1038/ni.2919
- Baruch, K., Rosenzweig, N., Kertser, A., Deczkowska, A., Sharif, A. M., Spinrad, A., Tsitsou-Kampeli, A., Sarel, A., Cahalon, L., & Schwartz, M. (2015). Breaking immune tolerance by targeting Foxp3(+) regulatory T cells mitigates Alzheimer's disease pathology. *Nature Communications*, 6, 7967. https://doi.org/10. 1038/ncomms8967
- Bergsbaken, T., Fink, S. L., & Cookson, B. T. (2009). Pyroptosis: Host cell death and inflammation. *Nature Reviews Microbiology*, 7(1), 99–109. https://doi.org/10.1038/ nrmicro2070
- Blacher, E., Dadali, T., Bespalko, A., Haupenthal, V. J., Grimm, M. O., Hartmann, T., Lund, F. E., Stein, R., & Levy, A. (2015). Alzheimer's disease pathology is attenuated in a CD38-deficient mouse model. *Annals of Neurology*, 78(1), 88–103. https://doi.org/10.1002/ana.24425
- Brazier, D., Perry, R., Keane, J., Barrett, K., & Elmaleh, D. R. (2017). Pharmacokinetics of cromolyn and ibuprofen in healthy elderly volunteers. *Clinical Drug Investigation*, 37, 1025–1034. https://doi.org/10.1007/s40261-017-0549-5
- Bsibsi, M., Ravid, R., Gveric, D., & Van Noort, J. M. (2002). Broad expression of toll-like receptors in the human central nervous system. *Journal of Neuropathology & Experimental Neurology*, 61(11), 1013–1021. https://doi.org/10.1093/jnen/ 61.11.1013
- Busche, M. A., & Hyman, B. T. (2020). Synergy between amyloid-β and tau in Alzheimer's disease. *Nature Neuroscience*, 23(10), 1183–1193. https://doi.org/10.1038/s41593-020-0687-6
- Caccamo, A., Branca, C., Piras, I. S., Ferreira, E., Huentelman, M. J., Liang, W. S., Readhead, B., Dudley, J. T., Spangenberg, E. E., Green, K. N., Belfiore, R., Winslow, W., & Oddo, S. (2017). Necroptosis activation in Alzheimer's disease. *Nature Neuroscience*, 20(9), 1236–1246. https://doi.org/10.1038/ nn.4608

- Cantrell, D. (2015). Signaling in lymphocyte activation. *Cold Spring Harbor Perspectives in Biology*, 7(6). https://doi.org/10.1101/ cshperspect.a018788
- Chasseigneaux, S., & Allinquant, B. (2012). Functions of Aβ, sAPPα and sAPPβ : Similarities and differences. *Journal of Neurochemistry*, *120*(Suppl 1), 99–108. https://doi.org/10.1111/ j.1471-4159.2011.07584.x
- Chen, G. F., Xu, T. H., Yan, Y., Zhou, Y. R., Jiang, Y., Melcher, K., & Xu, H. E. (2017). Amyloid beta: Structure, biology and structure-based therapeutic development. *Acta Pharmacologica Sinica*, 38(9), 1205–1235. https://doi.org/10.1038/aps.2017.28
- Chen, K., Iribarren, P., Hu, J., Chen, J., Gong, W., Cho, E. H., Lockett, S., Dunlop, N. M., & Wang, J. M. (2006). Activation of toll-like receptor 2 on microglia promotes cell uptake of Alzheimer disease-associated amyloid beta peptide. *Journal of Biological Chemistry*, 281(6), 3651–3659. https://doi.org/10. 1074/jbc.M508125200
- Christgen, S., Zheng, M., Kesavardhana, S., Karki, R., Malireddi, R. K. S., Banoth, B., Place, D. E., Briard, B., Sharma, B. R., Tuladhar, S., Samir, P., Burton, A., & Kanneganti, T. D. (2020). Identification of the PANoptosome: A molecular platform triggering pyroptosis, apoptosis, and necroptosis (PANoptosis). *Frontiers in Cellular and Infection Microbiology*, 10, 237. https://doi.org/10.3389/fcimb.2020.00237
- Clayton, K. A., Van Enoo, A. A., & Ikezu, T. (2017). Alzheimer's disease: The role of microglia in brain homeostasis and proteopathy. *Frontiers in Neuroscience*, 11, 680. https://doi.org/10.3389/ fnins.2017.00680
- Congdon, E. E., & Sigurdsson, E. M. (2018). Tau-targeting therapies for Alzheimer disease. *Nature Reviews. Neurology*, 14(7), 399– 415. https://doi.org/10.1038/s41582-018-0013-z
- Daniels, M. J., Rivers-Auty, J., Schilling, T., Spencer, N. G., Watremez, W., Fasolino, V., Booth, S. J., White, C. S., Baldwin, A. G., Freeman, S., Wong, R., Latta, C., Yu, S., Jackson, J., Fischer, N., Koziel, V., Pillot, T., Bagnall, J., Allan, S. M., ..., D. Brough (2016). Fenamate NSAIDs inhibit the NLRP3 inflammasome and protect against Alzheimer's disease in rodent models. *Nature Communications*, 7, 12504. https://doi.org/10.1038/ncomms12504
- Dansokho, C., Ait Ahmed, D., Aid, S., Toly-Ndour, C., Chaigneau, T., Calle, V., Cagnard, N., Holzenberger, M., Piaggio, E., Aucouturier, P., & Dorothée, G. (2016). Regulatory T cells delay disease progression in Alzheimer-like pathology. *Brain*, 139(Pt 4), 1237–1251. https://doi.org/10.1093/brain/awv408
- Decourt, B., Wilson, J., Ritter, A., Dardis, C., Difilippo, F. P., Zhuang, X., Cordes, D., Lee, G., Fulkerson, N. D., St Rose, T., Hartley, K., & Sabbagh, M. N. (2020). MCLENA-1: A phase II clinical trial for the assessment of safety, tolerability, and efficacy of lenalidomide in patients with mild cognitive impairment Due to Alzheimer's disease. *Open Access Journal of Clinical Trials*, *12*, 1–13. https://doi.org/10.2147/OAJCT.S221914
- De Rivero Vaccari, J. P., Brand, F. J.3rd, Sedaghat, C., Mash, D. C., Dietrich, W. D., & Keane, R. W. (2014). RIG-1 receptor expression in the pathology of Alzheimer's disease. *Journal of Neuroinflammation*, 11, 67. https://doi.org/10.1186/1742-2094-11-67
- Dondelinger, Y., Declercq, W., Montessuit, S., Roelandt, R., Goncalves, A., Bruggeman, I., Hulpiau, P., Weber, K., Sehon, C. A., Marquis, R. W., Bertin, J., Gough, P. J., Savvides, S., Martinou, J. C., Bertrand, M. J., & Vandenabeele, P. (2014).

MLKL Compromises plasma membrane integrity by binding to phosphatidylinositol phosphates. *Cell Reports*, 7(4), 971–981. https://doi.org/10.1016/j.celrep.2014.04.026

- Estus, S., Shaw, B. C., Devanney, N., Katsumata, Y., Press, E. E., & Fardo, D. W. (2019). Evaluation of CD33 as a genetic risk factor for Alzheimer's disease. *Acta Neuropathologica*, 138(2), 187– 199. https://doi.org/10.1007/s00401-019-02000-4
- Fiandaca, M. S., Kapogiannis, D., Mapstone, M., Boxer, A., Eitan, E., Schwartz, J. B., Abner, E. L., Petersen, R. C., Federoff, H. J., Miller, B. L., & Goetzl, E. J. (2015). Identification of preclinical Alzheimer's disease by a profile of pathogenic proteins in neurally derived blood exosomes: A case-control study. *Alzheimer's & Dementia*, 11(6), 600–607.e601. https://doi.org/ 10.1016/j.jalz.2014.06.0086
- Forlenza, O. V., Diniz, B. S., Talib, L. L., Mendonça, V. A., Ojopi, E. B., Gattaz, W. F., & Teixeira, A. L. (2009). Increased serum IL-1beta level in Alzheimer's disease and mild cognitive impairment. *Dementia and Geriatric Cognitive Disorders*, 28(6), 507– 512. https://doi.org/10.1159/000255051
- Forloni, G., Mangiarotti, F., Angeretti, N., Lucca, E., & De Simoni, M. G. (1997). Beta-amyloid fragment potentiates IL-6 and TNF-alpha secretion by LPS in astrocytes but not in microglia. *Cytokine*, 9(10), 759–762. https://doi.org/10.1006/cyto.1997. 0232
- Franklin, B. S., Bossaller, L., De Nardo, D., Ratter, J. M., Stutz, A., Engels, G., Brenker, C., Nordhoff, M., Mirandola, S. R., Al-Amoudi, A., Mangan, M. S., Zimmer, S., Monks, B. G., Fricke, M., Schmidt, R. E., Espevik, T., Jones, B., Jarnicki, A. G., Hansbro, P. M., ..., E. Latz (2014). The adaptor ASC has extracellular and 'prionoid' activities that propagate inflammation. *Nature Immunology*, 15(8), 727–737. https://doi.org/10. 1038/ni.2913
- Friker, L. L., Scheiblich, H., Hochheiser, I. V., Brinkschulte, R., Riedel, D., Latz, E., Geyer, M., & Heneka, M. T. (2020). β-Amyloid clustering around ASC fibrils boosts its toxicity in microglia. *Cell Reports*, 30(11), 3743–3754.e3746. https:// doi.org/10.1016/j.celrep.2020.02.025
- Gervais, F. G., Xu, D., Robertson, G. S., Vaillancourt, J. P., Zhu, Y., Huang, J., Leblanc, A., Smith, D., Rigby, M., Shearman, M. S., Clarke, E. E., Zheng, H., Van Der Ploeg, L. H., Ruffolo, S. C., Thornberry, N. A., Xanthoudakis, S., Zamboni, R. J., Roy, S., ... Nicholson, D. W. (1999). Involvement of caspases in proteolytic cleavage of Alzheimer's amyloid-beta precursor protein and amyloidogenic A beta peptide formation. *Cell*, 97(3), 395–406. https://doi.org/ 10.1016/S0092-8674(00)80748-5
- Giannakopoulos, P., Herrmann, F. R., Bussière, T., Bouras, C., Kövari, E., Perl, D. P., Morrison, J. H., Gold, G., & Hof, P. R. (2003). Tangle and neuron numbers, but not amyloid load, predict cognitive status in Alzheimer's disease. *Neurology*, 60(9), 1495–1500. https://doi.org/10.1212/01.WNL.0000063311.58879.01
- Go, M., Kou, J., Lim, J. E., Yang, J., & Fukuchi, K. I. (2016). Microglial response to LPS increases in wild-type mice during aging but diminishes in an Alzheimer's mouse model: Implication of TLR4 signaling in disease progression. *Biochemical and Biophysical Research Communications*, 479(2), 331–337. https://doi.org/10.1016/j.bbrc.2016.09.073
- Guerreiro, S., Privat, A. L., Bressac, L., & Toulorge, D. (2020). CD38 In neurodegeneration and neuroinflammation. *Cells*, 9(2). https://doi.org/10.3390/cells9020471

- Gurung, P., Anand, P. K., Malireddi, R. K., Vande Walle, L., Van Opdenbosch, N., Dillon, C. P., Weinlich, R., Green, D. R., Lamkanfi, M., & Kanneganti, T. D. (2014). FADD And caspase-8 mediate priming and activation of the canonical and noncanonical Nlrp3 inflammasomes. *Journal of Immunology*, 192(4), 1835–1846. https://doi.org/10.4049/jimmunol.1302839
- Gurung, P., & Kanneganti, T. D. (2016). Autoinflammatory skin disorders: The inflammasomme in focus. *Trends in Molecular Medicine*, 22(7), 545–564. https://doi.org/10.1016/j.molmed. 2016.05.003
- Haditsch, U., Roth, T., Rodriguez, L., Hancock, S., Cecere, T., Nguyen, M., Arastu-Kapur, S., Broce, S., Raha, D., Lynch, C. C., Holsinger, L. J., Dominy, S. S., & Ermini, F. (2020). Alzheimer's disease-like neurodegeneration in Porphyromonas gingivalis infected neurons with persistent expression of active gingipains. *Journal of Alzheimer's Disease*, 75(4), 1361–1376. https://doi.org/10.3233/JAD-200393
- Halle, A., Hornung, V., Petzold, G. C., Stewart, C. R., Monks, B. G., Reinheckel, T., Fitzgerald, K. A., Latz, E., Moore, K. J., & Golenbock, D. T. (2008). The NALP3 inflammasome is involved in the innate immune response to amyloid-beta. *Nature Immunology*, 9(8), 857–865. https://doi.org/10.1038/ni.1636
- Haller, O., Staeheli, P., Schwemmle, M., & Kochs, G. (2015). Mx GTPases: Dynamin-like antiviral machines of innate immunity. *Trends in Microbiology*, 23(3), 154–163. https://doi.org/10. 1016/j.tim.2014.12.003
- Han, C., Yang, Y., Guan, Q., Zhang, X., Shen, H., Sheng, Y., Wang, J., Zhou, X., Li, W., Guo, L., & Jiao, Q. (2020). New mechanism of nerve injury in Alzheimer's disease: B-amyloid-induced neuronal pyroptosis. *Journal of Cellular and Molecular Medicine*, 24(14), 8078–8090. https://doi.org/10.1111/jcmm.15439
- Hanisch, U. K., & Kettenmann, H. (2007). Microglia: Active sensor and versatile effector cells in the normal and pathologic brain. *Nature Neuroscience*, 10(11), 1387–1394. https://doi.org/10. 1038/nn1997
- Hansen, D. V., Hanson, J. E., & Sheng, M. (2018). Microglia in Alzheimer's disease. *Journal of Cell Biology*, 217(2), 459–472. https://doi.org/10.1083/jcb.201709069
- Hardy, J., & Selkoe, D. J. (2002). The amyloid hypothesis of Alzheimer's disease: Progress and problems on the road to therapeutics. *Science (new York, N Y)*, 297(2), 353–356. https://doi. org/10.1126/science.1072994
- He, W. T., Wan, H., Hu, L., Chen, P., Wang, X., Huang, Z., Yang, Z. H., Zhong, C. Q., & Han, J. (2015). Gasdermin D is an executor of pyroptosis and required for interleukin-1β secretion. *Cell Research*, 25(12), 1285–1298. https://doi.org/10.1038/cr.2015. 139
- He, Z., Guo, J. L., Mcbride, J. D., Narasimhan, S., Kim, H., Changolkar, L., Zhang, B., Gathagan, R. J., Yue, C., Dengler, C., Stieber, A., Nitla, M., Coulter, D. A., Abel, T., Brunden, K. R., Trojanowski, J. Q., & Lee, V. M. (2018). Amyloid-β plaques enhance Alzheimer's brain tau-seeded pathologies by facilitating neuritic plaque tau aggregation. *Nature Medicine*, 24(1), 29–38. https://doi.org/10.1038/nm.4443
- Hemonnot, A. L., Hua, J., Ulmann, L., & Hirbec, H. (2019). Microglia in Alzheimer disease: Well-known targets and new opportunities. *Frontiers in Aging Neuroscience*, 11, 233. https://doi.org/10.3389/fnagi.2019.00233
- Heneka, M. T., Carson, M. J., El Khoury, J., Landreth, G. E., Brosseron, F., Feinstein, D. L., Jacobs, A. H., Wyss-Coray, T.,

Vitorica, J., Ransohoff, R. M., Herrup, K., Frautschy, S. A., Finsen, B., Brown, G. C., Verkhratsky, A., Yamanaka, K., Koistinaho, J., Latz, E., Halle, A., ..., M. P. Kummer (2015). Neuroinflammation in Alzheimer's disease. *The Lancet. Neurology*, *14*(4), 388–405. https://doi.org/10.1016/S1474-4422(15)70016-5

- Heneka, M. T., Kummer, M. P., & Latz, E. (2014). Innate immune activation in neurodegenerative disease. *Nature Reviews Immunology*, 14(7), 463–477. https://doi.org/10.1038/nri3705
- Heneka, M. T., Kummer, M. P., Stutz, A., Delekate, A., Schwartz, S., Vieira-Saecker, A., Griep, A., Axt, D., Remus, A., Tzeng, T. C., Gelpi, E., Halle, A., Korte, M., Latz, E., & Golenbock, D. T. (2013). NLRP3 Is activated in Alzheimer's disease and contributes to pathology in APP/PS1 mice. *Nature*, 493(7434), 674– 678. https://doi.org/10.1038/nature11729
- Holmes, C., Boche, D., Wilkinson, D., Yadegarfar, G., Hopkins, V., Bayer, A., Jones, R. W., Bullock, R., Love, S., Neal, J. W., Zotova, E., & Nicoll, J. A. (2008). Long-term effects of Abeta42 immunisation in Alzheimer's disease: Follow-up of a randomised, placebo-controlled phase I trial. *Lancet (London, England)*, 372(9634), 216–223. https://doi.org/10.1016/S0140-6736(08)61075-2
- Ising, C., Venegas, C., Zhang, S., Scheiblich, H., Schmidt, S. V., Vieira-Saecker, A., Schwartz, S., Albasset, S., Mcmanus, R. M., Tejera, D., Griep, A., Santarelli, F., Brosseron, F., Opitz, S., Stunden, J., Merten, M., Kayed, R., Golenbock, D. T., & Blum, D., ..., M. T. Heneka (2019). NLRP3 Inflammasome activation drives tau pathology. *Nature*, 575(7784), 669–673. https://doi.org/10.1038/s41586-019-1769-z
- Jack, C. R. Jr., Knopman, D. S., Jagust, W. J., Shaw, L. M., Aisen, P. S., Weiner, M. W., Petersen, R. C., & Trojanowski, J. Q. (2010). Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *The Lancet. Neurology*, 9(1), 119–128. https://doi.org/10.1016/S1474-4422(09)70299-6
- Jana, M., Palencia, C. A., & Pahan, K. (2008). Fibrillar amyloid-beta peptides activate microglia via TLR2: Implications for Alzheimer's disease. *Journal of Immunology*, 181(10), 7254– 7262. https://doi.org/10.4049/jimmunol.181.10.7254
- Jia, L., Qiu, Q., Zhang, H., Chu, L., Du, Y., Zhang, J., Zhou, C., Liang, F., Shi, S., Wang, S., Qin, W., Wang, Q., Li, F., Wang, Q., Li, Y., Shen, L., Wei, Y., & Jia, J. (2019). Concordance between the assessment of Aβ42, T-tau, and P-T181-tau in peripheral blood neuronal-derived exosomes and cerebrospinal fluid. *Alzheimer's & Dementia*, 15(8), 1071–1080. https://doi. org/10.1016/j.jalz.2019.05.002
- Kanneganti, T. D. (2010). Central roles of NLRs and inflammasomes in viral infection. *Nature Reviews Immunology*, 10(10), 688–698. https://doi.org/10.1038/nri2851
- Kanneganti, T. D., Body-Malapel, M., Amer, A., Park, J. H., Whitfield, J., Franchi, L., Taraporewala, Z. F., Miller, D., Patton, J. T., Inohara, N., & Núñez, G. (2006a). Critical role for cryopyrin/Nalp3 in activation of caspase-1 in response to viral infection and double-stranded RNA. *Journal of Biological Chemistry*, 281(48), 36560–36568. https://doi.org/ 10.1074/jbc.M607594200
- Kanneganti, T. D., Ozören, N., Body-Malapel, M., Amer, A., Park, J. H., Franchi, L., Whitfield, J., Barchet, W., Colonna, M., Vandenabeele, P., Bertin, J., Coyle, A., Grant, E. P., Akira, S., & Núñez, G. (2006b). Bacterial RNA and small antiviral

compounds activate caspase-1 through cryopyrin/Nalp3. *Nature*, 440(7081), 233–236. https://doi.org/10.1038/ nature04517

- Karki, R., & Kanneganti, T. D. (2021). The 'cytokine storm': Molecular mechanisms and therapeutic prospects. *Trends in Immunology*, 42(8), 681–705. https://doi.org/10.1016/j.it.2021. 06.001
- Karki, R., Man, S. M., Malireddi, R. K. S., Gurung, P., Vogel, P., Lamkanfi, M., & Kanneganti, T. D. (2015). Concerted activation of the AIM2 and NLRP3 inflammasomes orchestrates host protection against Aspergillus infection. *Cell Host & Microbe*, *17*(3), 357–368. https://doi.org/10.1016/j.chom.2015.01.006
- Karki, R., Sharma, B. R., Tuladhar, S., Williams, E. P., Zalduondo, L., Samir, P., Zheng, M., Sundaram, B., Banoth, B., Malireddi, R. K. S., Schreiner, P., Neale, G., Vogel, P., Webby, R., Jonsson, C. B., & Kanneganti, T. D. (2021). Synergism of TNF-α and IFN-γ triggers inflammatory cell death, tissue damage, and mortality in SARS-CoV-2 infection and cytokine shock syndromes. *Cell*, *184*(1), 149–168.e117. https://doi.org/10.1016/j.cell.2020. 11.025
- Katzmarski, N., Ziegler-Waldkirch, S., Scheffler, N., Witt, C., Abou-Ajram, C., Nuscher, B., Prinz, M., Haass, C., & Meyer-Luehmann, M. (2020). Aβ oligomers trigger and accelerate Aβ seeding. *Brain Pathology*, 30(1), 36–45. https://doi.org/ 10.1111/bpa.12734
- Kesavardhana, S., Malireddi, R. K. S., Burton, A. R., Porter, S. N., Vogel, P., Pruett-Miller, S. M., & Kanneganti, T. D. (2020). The Zα2 domain of ZBP1 is a molecular switch regulating influenza-induced PANoptosis and perinatal lethality during development. *Journal of Biological Chemistry*, 295(24), 8325– 8330. https://doi.org/10.1074/jbc.RA120.013752
- Kigerl, K. A., De Rivero Vaccari, J. P., Dietrich, W. D., Popovich, P. G., & Keane, R. W. (2014). Pattern recognition receptors and central nervous system repair. *Experimental Neurology*, 258, 5–16. https://doi.org/10.1016/j.expneurol.2014.01.001
- King, E., O'brien, J. T., Donaghy, P., Morris, C., Barnett, N., Olsen, K., Martin-Ruiz, C., Taylor, J. P., & Thomas, A. J. (2018). Peripheral inflammation in prodromal Alzheimer's and Lewy body dementias. *Journal of Neurology Neurosurgery & Psychiatry*, 89(4), 339–345. https://doi.org/10.1136/jnnp-2017-317134
- Kuriakose, T., Man, S. M., Malireddi, R. K., Karki, R., Kesavardhana, S., Place, D. E., Neale, G., Vogel, P., & Kanneganti, T. D. (2016). ZBP1/DAI Is an innate sensor of influenza virus triggering the NLRP3 inflammasome and programmed cell death pathways. *Science Immunology*, 1(2). https://doi.org/ 10.1126/sciimmunol.aag2045
- Laferla, F. M., Tinkle, B. T., Bieberich, C. J., Haudenschild, C. C., & Jay, G. (1995). The Alzheimer's A beta peptide induces neurodegeneration and apoptotic cell death in transgenic mice. *Nature Genetics*, 9(1), 21–30. https://doi.org/10.1038/ng0195-21
- Laganke, C., Samkoff, L., Edwards, K., Jung Henson, L., Repovic, P., Lynch, S., Stone, L., Mattson, D., Galluzzi, A., Fisher, T. L., Reilly, C., Winter, L. A., Leonard, J. E., & Zauderer, M. (2017). Safety/tolerability of the anti-semaphorin 4D antibody VX15/2503 in a randomized phase 1 trial. *Neurology(R) Neuroimmunology & Neuroinflammation*, 4(4), e367. https:// doi.org/10.1212/NXI.00000000000367
- Lai, J., Mei, Z. L., Wang, H., Hu, M., Long, Y., Miao, M. X., Li, N., & Hong, H. (2014). Montelukast rescues primary neurons against

Aβ1-42-induced toxicity through inhibiting CysLT1R-mediated NF-κB signaling. *Neurochemistry International*, *75*, 26–31. https://doi.org/10.1016/j.neuint.2014.05.006

- Lamkanfi, M., Kanneganti, T. D., Van Damme, P., Vanden Berghe, T., Vanoverberghe, I., Vandekerckhove, J., Vandenabeele, P., Gevaert, K., & Núñez, G. (2008). Targeted peptidecentric proteomics reveals caspase-7 as a substrate of the caspase-1 inflammasomes. *Molecular and Cellular Proteomics*, 7(12), 2350–2363. https://doi.org/10.1074/mcp.M800132-MCP200
- Larbi, A., Pawelec, G., Witkowski, J. M., Schipper, H. M., Derhovanessian, E., Goldeck, D., & Fulop, T. (2009). Dramatic shifts in circulating CD4 but not CD8 T cell subsets in mild Alzheimer's disease. *Journal of Alzheimer's Disease*, 17(1), 91–103. https://doi.org/10.3233/JAD-2009-1015
- Lee, S., Channappanavar, R., & Kanneganti, T. D. (2020). Coronaviruses: Innate immunity, inflammasome activation, inflammatory cell death, and cytokines. *Trends in Immunology*, 41(12), 1083–1099. https://doi.org/10.1016/j.it. 2020.10.005
- Lee, S., Hirohama, M., Noguchi, M., Nagata, K., & Kawaguchi, A. (2018). Influenza A virus infection triggers pyroptosis and apoptosis of respiratory epithelial cells through the Type I interferon signaling pathway in a mutually exclusive manner. *Journal of Virology*, 92(14). https://doi.org/10.1128/jvi.00396-18
- Lee, S., Ishitsuka, A., Kuroki, T., Lin, Y. H., Shibuya, A., Hongu, T., Funakoshi, Y., Kanaho, Y., Nagata, K., & Kawaguchi, A. (2021a). Arf6 exacerbates allergic asthma through cell-to-cell transmission of ASC inflammasomes. *JCI Insight*, 6(16). https://doi.org/10.1172/jci.insight.139190
- Lee, S., Ishitsuka, A., Noguchi, M., Hirohama, M., Fujiyasu, Y., Petric, P. P., Schwemmle, M., Staeheli, P., Nagata, K., & Kawaguchi, A. (2019). Influenza restriction factor MxA functions as inflammasome sensor in the respiratory epithelium. *Science Immunology*, 4(40). https://doi.org/10.1126/ sciimmunol.aau4643
- Lee, S., Karki, R., Wang, Y., Nguyen, L. N., Kalathur, R. C., & Kanneganti, T. D. (2021b). AIM2 Forms a complex with pyrin and ZBP1 to drive PANoptosis and host defence. *Nature*, 597(7876), 415–419. https://doi.org/10.1038/s41586-021-03875-8
- Lee, S., & Ryu, J.-H. (2021). Influenza viruses: Innate immunity and mRNA vaccines. *Frontiers in Immunology*, 12(3534). https://doi. org/10.3389/fimmu.2021.710647
- Li, X., Melief, E., Postupna, N., Montine, K. S., Keene, C. D., & Montine, T. J. (2015). Prostaglandin E2 receptor subtype 2 regulation of scavenger receptor CD36 modulates microglial Aβ42 phagocytosis. *American Journal of Pathology*, 185(1), 230– 239. https://doi.org/10.1016/j.ajpath.2014.09.016
- Liu, S., Liu, Y., Hao, W., Wolf, L., Kiliaan, A. J., Penke, B., Rübe, C. E., Walter, J., Heneka, M. T., Hartmann, T., Menger, M. D., & Fassbender, K. (2012). TLR2 is a primary receptor for Alzheimer's amyloid β peptide to trigger neuroinflammatory activation. *Journal of Immunology*, *188*(3), 1098–1107. https://doi. org/10.4049/jimmunol.1101121
- Lonnemann, N., Hosseini, S., Marchetti, C., Skouras, D. B., Stefanoni, D., D'alessandro, A., Dinarello, C. A., & Korte, M. (2020). The NLRP3 inflammasome inhibitor OLT1177 rescues cognitive impairment in a mouse model of Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United*

States of America, 117(50), 32145–32154. https://doi.org/10. 1073/pnas.2009680117

- Lue, L. F., Walker, D. G., Brachova, L., Beach, T. G., Rogers, J., Schmidt, A. M., Stern, D. M., & Yan, S. D. (2001). Involvement of microglial receptor for advanced glycation endproducts (RAGE) in Alzheimer's disease: Identification of a cellular activation mechanism. *Experimental Neurology*, 171(1), 29–45. https://doi.org/10.1006/exnr.2001.7732
- Ma, S. L., Huang, W., Tang, N. L., & Lam, L. C. (2012). Mxa polymorphisms are associated with risk and age-at-onset in Alzheimer disease and accelerated cognitive decline in Chinese elders. *Rejuvenation Research*, 15(5), 516–522. https://doi.org/10.1089/rej.2012.1328
- Macpherson, K. P., Sompol, P., Kannarkat, G. T., Chang, J., Sniffen, L., Wildner, M. E., Norris, C. M., & Tansey, M. G. (2017).
 Peripheral administration of the soluble TNF inhibitor XPro1595 modifies brain immune cell profiles, decreases beta-amyloid plaque load, and rescues impaired long-term potentiation in 5xFAD mice. *Neurobiology of Disease*, *102*, 81–95. https://doi.org/10.1016/j.nbd.2017.02.010
- Malireddi, R. K., Ippagunta, S., Lamkanfi, M., & Kanneganti, T. D. (2010). Cutting edge: Proteolytic inactivation of poly(ADP-ribose) polymerase 1 by the Nlrp3 and Nlrc4 inflammasomes. *Journal of Immunology*, *185*(6), 3127–3130. https:// doi.org/10.4049/jimmunol.1001512
- Malireddi, R. K. S., Gurung, P., Kesavardhana, S., Samir, P., Burton, A., Mummareddy, H., Vogel, P., Pelletier, S., Burgula, S., & Kanneganti, T. D. (2020). Innate immune priming in the absence of TAK1 drives RIPK1 kinase activity-independent pyroptosis, apoptosis, necroptosis, and inflammatory disease. *Journal of Experimental Medicine*, 217(3). https://doi.org/10. 1084/jem.20191644
- Malireddi, R. K. S., Karki, R., Sundaram, B., Kancharana, B., Lee, S., Samir, P., & Kanneganti, T. D. (2021). Inflammatory cell death, PANoptosis, Mediated by cytokines in diverse cancer lineages inhibits tumor growth. *Immunohorizons*, 5(7), 568–580. https://doi.org/10.4049/immunohorizons.2100059
- Man, S. M., Karki, R., & Kanneganti, T. D. (2017). Molecular mechanisms and functions of pyroptosis, inflammatory caspases and inflammasomes in infectious diseases. *Immunological Reviews*, 277(1), 61–75. https://doi.org/10.1111/imr.12534
- Mancuso, R., Fryatt, G., Cleal, M., Obst, J., Pipi, E., Monzón-Sandoval, J., Ribe, E., Winchester, L., Webber, C., Nevado, A., Jacobs, T., Austin, N., Theunis, C., Grauwen, K., Daniela Ruiz, E., Mudher, A., Vicente-Rodriguez, M., Parker, C. A., Simmons, C., ..., V. H. Perry (2019). CSF1R Inhibitor JNJ-40346527 attenuates microglial proliferation and neurodegeneration in P301S mice. *Brain*, 142(10), 3243–3264. https:// doi.org/10.1093/brain/awz241
- Marsh, S. E., Abud, E. M., Lakatos, A., Karimzadeh, A., Yeung, S. T., Davtyan, H., Fote, G. M., Lau, L., Weinger, J. G., Lane, T. E., Inlay, M. A., Poon, W. W., & Blurton-Jones, M. (2016). The adaptive immune system restrains Alzheimer's disease pathogenesis by modulating microglial function. *Proceedings of the National Academy of Sciences of the United States of America*, 113(9), E1316–E1325. https://doi.org/10.1073/pnas.1525466113
- Mcdonald, C. L., Hennessy, E., Rubio-Araiz, A., Keogh, B., Mccormack, W., Mcguirk, P., Reilly, M., & Lynch, M. A. (2016). Inhibiting TLR2 activation attenuates amyloid accumulation and glial activation in a mouse model of Alzheimer's disease.

Brain Behavior and Immunity, 58, 191–200. https://doi.org/10. 1016/j.bbi.2016.07.143

- Mcdonough, A., Lee, R. V., & Weinstein, J. R. (2017). Microglial interferon signaling and white matter. *Neurochemical Research*, 42(9), 2625–2638. https://doi.org/10.1007/s11064-017-2307-8
- Morin, N., Jourdain, V. A., Morissette, M., Grégoire, L., & Di Paolo, T. (2014). Long-term treatment with 1-DOPA and an mGlu5 receptor antagonist prevents changes in brain basal ganglia dopamine receptors, their associated signaling proteins and neuropeptides in parkinsonian monkeys. *Neuropharmacology*, 79, 688–706. https://doi.org/10.1016/j.neuropharm.2014.01.014
- Nelson, P. T., Alafuzoff, I., Bigio, E. H., Bouras, C., Braak, H., Cairns, N. J., Castellani, R. J., Crain, B. J., Davies, P., Del Tredici, K., Duyckaerts, C., Frosch, M. P., Haroutunian, V., Hof, P. R., Hulette, C. M., Hyman, B. T., Iwatsubo, T., Jellinger, K. A., Jicha, G. A., ..., T. G. Beach (2012). Correlation of Alzheimer disease neuropathologic changes with cognitive status: A review of the literature. *Journal of Neuropathology & Experimental Neurology*, *71*(5), 362–381. https://doi.org/10.1097/NEN.0b013e31825018f7
- Ng, A., Tam, W. W., Zhang, M. W., Ho, C. S., Husain, S. F., Mcintyre, R. S., & Ho, R. C. (2018). IL-1β, IL-6, TNF- α and CRP in elderly patients with depression or Alzheimer's disease: Systematic review and meta-analysis. *Scientific Reports*, 8(1), 12050. https://doi.org/10.1038/s41598-018-30487-6
- Nussbaum, R. L., & Ellis, C. E. (2003). Alzheimer's disease and Parkinson's disease. New England Journal of Medicine, 348(14), 1356–1364. https://doi.org/10.1056/ NEJM2003ra020003
- Ofengeim, D., Mazzitelli, S., Ito, Y., Dewitt, J. P., Mifflin, L., Zou, C., Das, S., Adiconis, X., Chen, H., Zhu, H., Kelliher, M. A., Levin, J. Z., & Yuan, J. (2017). RIPK1 Mediates a disease-associated microglial response in Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America*, 114(41), E8788–e8797. https://doi.org/10. 1073/pnas.1714175114
- Ojala, J., Alafuzoff, I., Herukka, S. K., Van Groen, T., Tanila, H., & Pirttilä, T. (2009). Expression of interleukin-18 is increased in the brains of Alzheimer's disease patients. *Neurobiology of Aging*, 30(2), 198–209. https://doi.org/10.1016/j.neurobiolaging.2007. 06.006
- Olson, J. K., & Miller, S. D. (2004). Microglia initiate central nervous system innate and adaptive immune responses through multiple TLRs. *Journal of Immunology*, 173(6), 3916–3924. https://doi.org/10.4049/jimmunol.173.6.3916
- Palm, N. W., & Medzhitov, R. (2009). Pattern recognition receptors and control of adaptive immunity. *Immunological Reviews*, 227(1), 221–233. https://doi.org/10.1111/j.1600-065X.2008. 00731.x
- Pasparakis, M., & Vandenabeele, P. (2015). Necroptosis and its role in inflammation. *Nature*, 517(7534), 311–320. https://doi.org/10. 1038/nature14191
- Pellicanò, M., Larbi, A., Goldeck, D., Colonna-Romano, G., Buffa, S., Bulati, M., Rubino, G., Iemolo, F., Candore, G., Caruso, C., Derhovanessian, E., & Pawelec, G. (2012). Immune profiling of Alzheimer patients. *Journal of Neuroimmunology*, 242(1-2), 52–59. https://doi.org/10.1016/j.jneuroim.2011.11.005
- Pirttilä, T., Mattinen, S., & Frey, H. (1992). The decrease of CD8-positive lymphocytes in Alzheimer's disease. *Journal of*

the Neurological Sciences, *107*(2), 160–165. https://doi.org/10. 1016/0022-510X(92)90284-R

- Place, D. E., & Kanneganti, T. D. (2018). Recent advances in inflammasome biology. *Current Opinion in Immunology*, 50, 32–38. https://doi.org/10.1016/j.coi.2017.10.011
- Place, D. E., Lee, S., & Kanneganti, T. D. (2021). PANoptosis in microbial infection. *Current Opinion in Microbiology*, 59, 42–49. https://doi.org/10.1016/j.mib.2020.07.012
- Plascencia-Villa, G., & Perry, G. (2020). Status and future directions of clinical trials in Alzheimer's disease. *International Review of Neurobiology*, 154, 3–50. https://doi.org/10.1016/bs.irn.2020. 03.022
- Qin, Y., Liu, Y., Hao, W., Decker, Y., Tomic, I., Menger, M. D., Liu, C., & Fassbender, K. (2016). Stimulation of TLR4 attenuates Alzheimer's disease-related symptoms and pathology in tautransgenic mice. *Journal of Immunology*, 197(8), 3281–3292. https://doi.org/10.4049/jimmunol.1600873
- Reed-Geaghan, E. G., Savage, J. C., Hise, A. G., & Landreth, G. E. (2009). CD14 And toll-like receptors 2 and 4 are required for fibrillar A{beta}-stimulated microglial activation. *Journal of Neuroscience*, 29(38), 11982–11992. https://doi.org/10.1523/ JNEUROSCI.3158-09.2009
- Rogers, C., Erkes, D. A., Nardone, A., Aplin, A. E., Fernandes-Alnemri, T., & Alnemri, E. S. (2019). Gasdermin pores permeabilize mitochondria to augment caspase-3 activation during apoptosis and inflammasome activation. *Nature Communications*, 10(1), 1689. https://doi.org/10.1038/s41467-019-09397-2
- Rohn, T. T., Head, E., Nesse, W. H., Cotman, C. W., & Cribbs, D. H. (2001). Activation of caspase-8 in the Alzheimer's disease brain. *Neurobiology of Disease*, 8(6), 1006–1016. https://doi.org/10. 1006/nbdi.2001.0449
- Rohn, T. T., Rissman, R. A., Davis, M. C., Kim, Y. E., Cotman, C. W., & Head, E. (2002). Caspase-9 activation and caspase cleavage of tau in the Alzheimer's disease brain. *Neurobiology of Disease*, *11*(2), 341–354. https://doi.org/10.1006/nbdi.2002.0549
- Roy, E. R., Wang, B., Wan, Y. W., Chiu, G., Cole, A., Yin, Z., Propson, N. E., Xu, Y., Jankowsky, J. L., Liu, Z., Lee, V. M., Trojanowski, J. Q., Ginsberg, S. D., Butovsky, O., Zheng, H., & Cao, W. (2020). Type I interferon response drives neuroinflammation and synapse loss in Alzheimer disease. *Journal of Clinical Investigation*, *130*(4), 1912–1930. https://doi.org/10.1172/ JCI133737
- Rubio-Perez, J. M., & Morillas-Ruiz, J. M. (2012). A review: Inflammatory process in Alzheimer's disease, role of cytokines. *Thescientificworldjournal*, 2012, 756357. 10.1100/ 2012/756357
- Salter, M. W., & Stevens, B. (2017). Microglia emerge as central players in brain disease. *Nature Medicine*, 23(9), 1018–1027. https://doi.org/10.1038/nm.4397
- Saresella, M., Calabrese, E., Marventano, I., Piancone, F., Gatti, A., Alberoni, M., Nemni, R., & Clerici, M. (2011). Increased activity of Th-17 and Th-9 lymphocytes and a skewing of the post-thymic differentiation pathway are seen in Alzheimer's disease. *Brain Behavior and Immunity*, 25(3), 539–547. https://doi.org/10. 1016/j.bbi.2010.12.004
- Saresella, M., La Rosa, F., Piancone, F., Zoppis, M., Marventano, I., Calabrese, E., Rainone, V., Nemni, R., Mancuso, R., & Clerici, M. (2016). The NLRP3 and NLRP1 inflammasomes are activated

in Alzheimer's disease. *Molecular Neurodegeneration*, 11, 23. https://doi.org/10.1186/s13024-016-0088-1

- Sarlus, H., & Heneka, M. T. (2017). Microglia in Alzheimer's disease. *Journal of Clinical Investigation*, 127(9), 3240–3249. https://doi.org/10.1172/JCI90606
- Schindowski, K., Peters, J., Gorriz, C., Schramm, U., Weinandi, T., Leutner, S., Maurer, K., Frölich, L., Müller, W. E., & Eckert, A. (2006). Apoptosis of CD4+ T and natural killer cells in Alzheimer's disease. *Pharmacopsychiatry*, 39(6), 220–228. https://doi.org/10.1055/s-2006-954591
- Selkoe, D. J., & Hardy, J. (2016). The amyloid hypothesis of Alzheimer's disease at 25 years. *Embo Molecular Medicine*, 8(6), 595–608. https://doi.org/10.15252/emmm.201606210
- Sevigny, J., Chiao, P., Bussière, T., Weinreb, P. H., Williams, L., Maier, M., Dunstan, R., Salloway, S., Chen, T., Ling, Y., O'gorman, J., Qian, F., Arastu, M., Li, M., Chollate, S., Brennan, M. S., Quintero-Monzon, O., Scannevin, R. H., Arnold, H. M., ..., A. Sandrock (2016). The antibody aducanumab reduces Aβ plaques in Alzheimer's disease. *Nature*, 537(7618), 50–56. https://doi.org/10.1038/nature19323
- Shi, J., Zhao, Y., Wang, K., Shi, X., Wang, Y., Huang, H., Zhuang, Y., Cai, T., Wang, F., & Shao, F. (2015). Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature*, 526(7575), 660–665. https://doi.org/10.1038/nature15514
- Song, M., Jin, J., Lim, J. E., Kou, J., Pattanayak, A., Rehman, J. A., Kim, H. D., Tahara, K., Lalonde, R., & Fukuchi, K. (2011). TLR4 Mutation reduces microglial activation, increases Aβ deposits and exacerbates cognitive deficits in a mouse model of Alzheimer's disease. *Journal of Neuroinflammation*, 8, 92. https://doi.org/ 10.1186/1742-2094-8-92
- Speciale, L., Calabrese, E., Saresella, M., Tinelli, C., Mariani, C., Sanvito, L., Longhi, R., & Ferrante, P. (2007). Lymphocyte subset patterns and cytokine production in Alzheimer's disease patients. *Neurobiology of Aging*, 28(8), 1163–1169. https://doi. org/10.1016/j.neurobiolaging.2006.05.020
- Stadelmann, C., Deckwerth, T. L., Srinivasan, A., Bancher, C., Brück, W., Jellinger, K., & Lassmann, H. (1999). Activation of caspase-3 in single neurons and autophagic granules of granulovacuolar degeneration in Alzheimer's disease. Evidence for apoptotic cell death. *American Journal of Pathology*, 155(5), 1459–1466. https://doi.org/10.1016/S0002-9440(10) 65460-0
- St-Amour, I., Paré, I., Tremblay, C., Coulombe, K., Bazin, R., & Calon, F. (2014). IVIg protects the 3xTg-AD mouse model of Alzheimer's disease from memory deficit and Aβ pathology. *Journal of Neuroinflammation*, 11, 54. https://doi.org/10.1186/ 1742-2094-11-54
- Stancu, I. C., Cremers, N., Vanrusselt, H., Couturier, J., Vanoosthuyse, A., Kessels, S., Lodder, C., Brône, B., Huaux, F., Octave, J. N., Terwel, D., & Dewachter, I. (2019). Aggregated Tau activates NLRP3-ASC inflammasome exacerbating exogenously seeded and non-exogenously seeded Tau pathology in vivo. Acta Neuropathologica, 137(4), 599–617. https://doi.org/10.1007/s00401-018-01957-y
- Stienstra, R., Van Diepen, J. A., Tack, C. J., Zaki, M. H., Van De Veerdonk, F. L., Perera, D., Neale, G. A., Hooiveld, G. J., Hijmans, A., Vroegrijk, I., Van Den Berg, S., Romijn, J., Rensen, P. C., Joosten, L. A., Netea, M. G., & Kanneganti, T. D. (2011). Inflammasome is a central player in the induction of obesity and insulin resistance. *Proceedings of the National*

Academy of Sciences of the United States of America, 108(37), 15324–15329. https://doi.org/10.1073/pnas.1100255108

- Su, J. H., Zhao, M., Anderson, A. J., Srinivasan, A., & Cotman, C. W. (2001). Activated caspase-3 expression in Alzheimer's and aged control brain: Correlation with Alzheimer pathology. *Brain Research*, 898(2), 350–357. https://doi.org/10.1016/ S0006-8993(01)02018-2
- Tahara, K., Kim, H. D., Jin, J. J., Maxwell, J. A., Li, L., & Fukuchi, K. (2006). Role of toll-like receptor signalling in Abeta uptake and clearance. *Brain*, 129(Pt 11), 3006–3019. https://doi.org/10. 1093/brain/awl249
- Tan, M. S., Tan, L., Jiang, T., Zhu, X. C., Wang, H. F., Jia, C. D., & Yu, J. T. (2014). Amyloid-β induces NLRP1-dependent neuronal pyroptosis in models of Alzheimer's disease. *Cell Death & Disease*, 5(8), e1382. https://doi.org/10.1038/cddis.2014.348
- Thomas, P. G., Dash, P., Aldridge, J. R. Jr., Ellebedy, A. H., Reynolds, C., Funk, A. J., Martin, W. J., Lamkanfi, M., Webby, R. J., Boyd, K. L., Doherty, P. C., & Kanneganti, T. D. (2009). The intracellular sensor NLRP3 mediates key innate and healing responses to influenza A virus via the regulation of caspase-1. *Immunity*, 30(4), 566–575. https://doi.org/10. 1016/j.immuni.2009.02.006
- Venegas, C., Kumar, S., Franklin, B. S., Dierkes, T., Brinkschulte, R., Tejera, D., Vieira-Saecker, A., Schwartz, S., Santarelli, F., Kummer, M. P., Griep, A., Gelpi, E., Beilharz, M., Riedel, D., Golenbock, D. T., Geyer, M., Walter, J., Latz, E., Heneka, M. T., ... (2017). Microglia-derived ASC specks cross-seed amyloid-β in Alzheimer's disease. *Nature*, 552(7685), 355–361. https://doi.org/10.1038/nature25158
- Vollmar, P., Kullmann, J. S., Thilo, B., Claussen, M. C., Rothhammer, V., Jacobi, H., Sellner, J., Nessler, S., Korn, T., & Hemmer, B. (2010). Active immunization with amyloid-beta 1-42 impairs memory performance through TLR2/4-dependent activation of the innate immune system. *Journal of Immunology*, 185(10), 6338–6347. https://doi.org/10.4049/ jimmunol.1001765
- Walter, S., Letiembre, M., Liu, Y., Heine, H., Penke, B., Hao, W., Bode, B., Manietta, N., Walter, J., Schulz-Schuffer, W., & Fassbender, K. (2007). Role of the toll-like receptor 4 in neuroinflammation in Alzheimer's disease. *Cellular Physiology and Biochemistry*, 20(6), 947–956. https://doi.org/10.1159/ 000110455
- Wang, S., Mustafa, M., Yuede, C. M., Salazar, S. V., Kong, P., Long, H., Ward, M., Siddiqui, O., Paul, R., Gilfillan, S., Ibrahim, A., Rhinn, H., Tassi, I., Rosenthal, A., Schwabe, T., & Colonna, M. (2020). Anti-human TREM2 induces microglia proliferation and reduces pathology in an Alzheimer's disease model. *Journal of Experimental Medicine*, 217(9). https://doi.org/10. 1084/jem.20200785
- Wang, W. Y., Tan, M. S., Yu, J. T., & Tan, L. (2015). Role of pro-inflammatory cytokines released from microglia in Alzheimer's disease. *Annals of Translational Medicine*, 3(10), 136. https://doi.org/10.3978/j.issn.2305-5839.2015.03.49
- Wesseling, H., Mair, W., Kumar, M., Schlaffner, C. N., Tang, S., Beerepoot, P., Fatou, B., Guise, A. J., Cheng, L., Takeda, S., Muntel, J., Rotunno, M. S., Dujardin, S., Davies, P., Kosik, K. S., Miller, B. L., Berretta, S., Hedreen, J. C., Grinberg, L. T., ..., J. A. Steen (2020). Tau PTM profiles identify patient heterogeneity and stages of Alzheimer's disease. *Cell*, 183(6), 1699– 1713.e1613. https://doi.org/10.1016/j.cell.2020.10.029

- Wu, P. J., Hung, Y. F., Liu, H. Y., & Hsueh, Y. P. (2017). Deletion of the inflammasome sensor Aim2 mitigates Aβ deposition and microglial activation but increases inflammatory cytokine expression in an Alzheimer disease mouse model. *Neuroimmunomodulation*, 24(1), 29–39. https://doi.org/10.1159/000477092
- Xia, M. Q., & Hyman, B. T. (1999). Chemokines/chemokine receptors in the central nervous system and Alzheimer's disease. *Journal of Neurovirology*, 5(1), 32–41. https://doi.org/10.3109/ 13550289909029743
- Yamada, T., Horisberger, M. A., Kawaguchi, N., Moroo, I., & Toyoda, T. (1994). Immunohistochemistry using antibodies to alpha-interferon and its induced protein, MxA, in Alzheimer's and Parkinson's disease brain tissues. *Neuroscience Letters*, 181(1-2), 61–64. https://doi.org/10. 1016/0304-3940(94)90560-6
- Yin, J., Zhao, F., Chojnacki, J. E., Fulp, J., Klein, W. L., Zhang, S., & Zhu, X. (2018). NLRP3 Inflammasome inhibitor ameliorates amyloid pathology in a mouse model of Alzheimer's disease. *Molecular Neurobiology*, 55(3), 1977–1987. https://doi.org/10. 1007/s12035-017-0467-9
- Zaki, M. H., Boyd, K. L., Vogel, P., Kastan, M. B., Lamkanfi, M., & Kanneganti, T. D. (2010). The NLRP3 inflammasome protects against loss of epithelial integrity and mortality during experimental colitis. *Immunity*, 32(3), 379–391. https://doi.org/10. 1016/j.immuni.2010.03.003
- Zenaro, E., Pietronigro, E., Della Bianca, V., Piacentino, G., Marongiu, L., Budui, S., Turano, E., Rossi, B., Angiari, S., Dusi, S., Montresor, A., Carlucci, T., Nanì, S., Tosadori, G., Calciano, L., Catalucci, D., Berton, G., Bonetti, B., Constantin, G., ... (2015). Neutrophils promote Alzheimer's disease-like pathology and cognitive decline via LFA-1 integrin. *Nature Medicine*, 21(8), 880–886. https://doi.org/10.1038/nm.3913
- Zhang, C., Griciuc, A., Hudry, E., Wan, Y., Quinti, L., Ward, J., Forte, A. M., Shen, X., Ran, C., Elmaleh, D. R., & Tanzi, R. E. (2018). Cromolyn reduces levels of the Alzheimer's disease-associated amyloid β-protein by promoting microglial phagocytosis. *Scientific Reports*, 8(1), 1144. https://doi.org/10. 1038/s41598-018-19641-2
- Zhang, P., Kishimoto, Y., Grammatikakis, I., Gottimukkala, K., Cutler, R. G., Zhang, S., Abdelmohsen, K., Bohr, V. A., Misra

Sen, J., Gorospe, M., & Mattson, M. P. (2019). Senolytic therapy alleviates A β -associated oligodendrocyte progenitor cell senescence and cognitive deficits in an Alzheimer's disease model. *Nature Neuroscience*, 22(5), 719–728. https://doi.org/10.1038/s41593-019-0372-9

- Zhang, Y. K., Liu, J. T., Peng, Z. W., Fan, H., Yao, A. H., Cheng, P., Liu, L., Ju, G., & Kuang, F. (2013). Different TLR4 expression and microglia/macrophage activation induced by hemorrhage in the rat spinal cord after compressive injury. *Journal of Neuroinflammation*, 10, 112. https://doi.org/10.1186/1742-2094-10-112
- Zheng, M., Karki, R., Vogel, P., & Kanneganti, T. D. (2020). Caspase-6 is a key regulator of innate immunity, inflammasome activation, and host defense. *Cell*, 181(3), 674–687.e613. https://doi.org/10.1016/j.cell.2020.03.040

Abbreviations

AD	Alzheimer's disease
APP	amyloid precursor protein
ASC	apoptosis-associated speck-like protein
	containing a CARD
Αβ	β-amyloid
DAMP	danger-associated molecular pattern
GSDMD	N-terminal fragment of gasdermin D
IFN	interferon
IL	interleukin
MCP-1	monocyte chemoattractant protein-1
MLKL	mixed-lineage kinase domain-like pseudokinase
NLR	nucleotide-binding oligomerization domain-like
	receptor family protein
NLRP3	NLR family pyrin domain-containing 3
PAMP	pathogen-associated molecular pattern
PRR	pattern-recognition receptor
PTM	post-translational modification
RIG-I	retinoic acid-inducible gene I
RLR	retinoic acid-inducible gene I-like receptor
TLR	Toll-like receptor.